

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Perfluoroalkyl Substances, Sex Hormones, and Insulin-like Growth Factor-1 at 6–9 Years of Age: A Cross-Sectional Analysis within the C8 Health Project

Maria-Jose Lopez-Espinosa, Debapriya Mondal, Ben G. Armstrong, Brenda Eskenazi, and Tony Fletcher

Table of Contents

Table S1. Number (%) of imputed values and regression model for each variable (*n*: 2,292 cases), Mid-Ohio Valley (USA), 2005-2006

Table S2. Pearson's correlations between ln(PFAS) in children aged 6-9 years, Mid-Ohio Valley (USA), 2005-2006

Table S3. Difference in levels of sex hormones and IGF-1 in relation to PFAS in boys aged 6-9 years, Mid-Ohio Valley (USA), 2005-2006

Table S4. Difference in levels of sex hormones and IGF-1 in relation to PFAS in girls aged 6-9 years, Mid-Ohio Valley (USA), 2005-2006

Table S5. Difference in levels of sex hormones and IGF-1 by PFAS quartiles in boys aged 6-9 years, Mid-Ohio Valley (USA), 2005-2006

Table S6. Difference in levels of sex hormones and IGF-1 by PFAS quartiles in girls aged 6-9 years, Mid-Ohio Valley (USA), 2005-2006

References